
# ATHENA.

#### Athena: ESA's X-ray observatory to study the Hot and Energetic Universe





# Matteo Guainazzi on behalf of the Athena Science Study Team


ESA/SCI-S, ESTEC, Noordwijk



#### Advanced Telescope for High-Energy Astrophysics

- Second Large (L2) mission of ESA Cosmic Vision 2015-2035
  - International contribution by JAXA and NASA
- Science theme: The Hot and Energetic Universe
  - How does ordinary matter assemble in the large-scale structures?
  - How do black holes grow and shape galaxies?
- In addition:
  - ToO capability to study transient sources
  - Observatory science across all corners of Astrophysics

More info at: <u>http://www.the-athena-x-ray-observatory.eu</u>




Athena widelv discussed at the

800+ scientists in the Athena community



# Mission profile (current Phase A)






- Single telescope, Silicon Pore Optics (SPO) technology, 12 m focal length, ≥1.4 m<sup>2</sup> area@1 keV, 0.25 m<sup>2</sup> @6 keV
- WFI (Active Pixel Sensor Si detector): wide-field (40'x40') spectral-imaging, CCD-like energy resolution (120-150 eV @6 keV)
- X-IFU (cryogenic imaging spectrometer): 2.5 eV energy resolution, 5'x5' field-of-view, ~5" pixel size
- Count rates capabilities: >1 Crab (WFI)/~1 Crab (30% throughput) X-IFU (increased thanks to defocusing capabilities)
- ≥4 hours response with a ~50% efficiency to observe a ToO in a random position in the sky
- Launch 2028/9, Ariane 6.4, L2 halo orbit (TBC)
- Nominal life-time 4 years + extensions

#### A revolutionary mix of science performance - I


Effective area per energy resolution element

Simulated velocity map at a 5" pixel resolution






#### A revolutionary mix of science performance - II





### Study status




Next key date: 2020, adoption (= final inclusion in the ESA Science program)



#### **Optics status**

May 2016 measurements at BESSY Half Energy Width per column 20 m focal length optics



- <HEW>: ~22" in 2015 → 13.9" in 2016
- 60% of the optics have a HEW of 8"
- Best performance: ~5"
- Consistent results at BESSY (2.8 keV) and Panter (1.49 keV)
- Next measurement campaign (with 12 m focal length optics): fall 2017





### WFI status

- Prototype sensor produced
  - 64x64 pixel matrix, 2.0-2.5 e<sup>-</sup> r.m.s., FWHM≈130 eV
  - 256x256 pixel matrix, 2.5 e<sup>-</sup> r.m.s, FWHM≈134 eV

# of events

- Detector electronics frame processor module for real-time event processing at 100 Mpixel/s set-up for testing
- Instrument design w/o vacuum enclosure for the Optical Blocking Filter successfully tested based on Ariane-5 launch reference









### X-IFU status

- 50 mK cooling chain technology demonstrator being developed under ESA contract with CNES lead and X-IFU consortium partners
- Large format TES arrays being fabricated and tested at GSFC
- Frequency domain multiplexing approach reaching the required resolution
- New (and promising) cryo-chain architecture under analysis. First results on the thermal budget expected in the next months











#### The Hot Universe - I.

#### Chemical evolution of the inter-cluster gas

- Clusters of galaxies are closed boxes, all gas is virialised in the DM potential well
- Cosmic chemical evolution best traced by cluster gas
- Constraints on SN types and IMF
- Probing clusters and groups up to z~2








#### The Hot Universe - II.


### AGN feedback on cluster scales

#### Dissipation AGN energy into ICM

- Energy stored in hot gas around bubbles via bulk motions and turbulence.
- History of radio cluster feedback via ripples.
- AGN jet fuelling vs. cooling through temperature distribution.
- Shock speeds of expanding radio lobes



A. Keesa




#### The Hot Universe - II.


### AGN feedback on cluster scales

#### Dissipation AGN energy into ICM

- Energy stored in hot gas around bubbles via bulk motions and turbulence.
- History of radio cluster feedback via ripples.
- AGN jet fuelling vs. cooling through temperature distribution.
- Shock speeds of expanding radio lobes



A. Keesa




#### The Hot Universe - III.

# Missing baryons: the WHIM\* \* Warm-Hot Intergalactic Medium

- Cosmological hydro simulations show ~50% of baryons at **T~10<sup>5</sup>-10**<sup>7</sup> K in the IGM.
  - Unvirialised and filamentary distribution
- How can they be detected?
  - In absorption:
    - Against a bright background sources
  - In emission:
    - Tenuous and extended
    - Key to understand CGM and feedback






#### The Energetic Universe - I.

#### AGN disk wind feedback with Athena

- AGN outflows with L<sub>mech</sub>≥1%L<sub>bol</sub> may be the "feedback messenger"
- Relativistic (v≥0.1c) disk outflows discovered at X-ray CCD-resolution However:
  - no plasma diagnostic possible
  - no estimate of mass and kinetic energy outflow possible
- High-resolution at the Fe band (6-7 keV) is the key
- Athena will make this possible, up to z~4






The Energetic Universe - II.

#### Obscured AGN census @ z~1-3


#AthenaNuggets Carrera

- Most SMBH growth expected in heavily obscured environment.
- Athena/WFI observations can uncover Compton-Thick (N<sub>H</sub>≥10<sup>24</sup> cm<sup>-2</sup>) AGN at z~3
  - MIR observations can reliably uncover heavily obscured AGN, but only when the AGN is very powerful
- Expected about 60 Comptonthick AGN ( $1 \le z \le 3$ ) over 6 degrees<sup>2</sup> down to  $L_x \sim 10^{44}$  erg s<sup>-1</sup>



The Energetic Universe - III.


#### The history of SMBH growth



#### Observatory Science – all corners of astrophysics

- Planets and solar system bodies
- Star formation, brown dwarfs
- Massive stars: mass loss
- Outflows in X-ray binaries
- Supernovae: explosion mechanisms
- Supernova remnants: shock physics
- Interstellar medium
- Dark Matter candidates

Cesa







- Athena will be a transformational X-ray observatory
  - Designed to address the Hot and Energetic Universe science theme
  - Will impact virtually every corner of astronomy



- Athena will be a transformational X-ray observatory
  - Designed to address the Hot and Energetic Universe science theme
  - Will impact virtually every corner of astronomy
- It will be an essential part of the observational landscape in the late 2020s, together with ALMA, E-ELT, SKA, CTA, etc.



- Athena will be a transformational X-ray observatory
  - Designed to address the Hot and Energetic Universe science theme
  - Will impact virtually every corner of astronomy
- It will be an essential part of the observational landscape in the late 2020s, together with ALMA, E-ELT, SKA, CTA, etc.
- Vibrant community supporting it



- Athena will be a transformational X-ray observatory
  - Designed to address the Hot and Energetic Universe science theme
  - Will impact virtually every corner of astronomy
- It will be an essential part of the observational landscape in the late 2020s, together with ALMA, E-ELT, SKA, CTA, etc.
- Vibrant community supporting it
- Good progress with Phase A.
  - Key milestone in 2020: Mission adoption by ESA for a launch in 2028/9.



- Athena will be a transformational X-ray observatory
  - Designed to address the Hot and Energetic Universe science theme
  - Will impact virtually every corner of astronomy
- It will be an essential part of the observational landscape in the late 2020s, together with ALMA, E-ELT, SKA, CTA, etc.
- Vibrant community supporting it
- Good progress with Phase A.
  - Key milestone in 2020: Mission adoption by ESA for a launch in 2028/9.
    - Follow Athena on
      - Web: www.the-athena-x-ray-observatory.eu
      - Twitter: @athena2028
      - Facebook: The Athena X-ray Observatory
      - Athena Community Office email: <u>aco@ifca.unican.es</u>

